

PSJ17 Exh 94

**From:** Catherine Collier  
**To:** Castagno, Paula  
**CC:** Beckhardt, Stacey  
**Sent:** 10/2/2006 7:55:38 PM  
**Subject:** ACPA brochure, etc  
**Attachments:** Breakthrough Pain 9-20-06.pdf

Hi Paula,

Hope you had an uneventful and safe flight back to Philly. I drove Stacey to airport at about 85 miles per hour to assure that she did not miss her flight ....think we cut it a bit short but the good news is we made it safely and she caught her flight.

In any event, as a follow up to our conversation on Friday, attached please find a copy of the ACPA brochure on BTP that Cephalon sponsored as part of the September Pain Awareness Month activities. This will soon be posted on their website homepage under the "News" section. In addition, they will be including it in every informational packet they send out. Once we have hard copy I will send you a few for your archives.

ACPA is also hosting a radio media tour tomorrow on breakthrough pain with Penney Cowan as the person with pain, and Dr. Knox Todd, Professor of Emergency Medicine

Director at Beth Israel Medical Center. If you are interested I can send you additional information on this as well.

Finally, I have an invoice for the \$60K that we discussed for activities related to launch (media montage, additional b-roll). I will fax it to you later today; please let me know if you need additional details added (e.g. budget estimates).

Thanks Paula,

Catherine



### What Causes It?

BTP can hit unexpectedly at any time or place and it may be hard to identify what triggers it. A cough or rolling over in bed can cause BTP. Or it can happen when a dose of your persistent pain medicine wears off before it is time to take another dose.

### The Impact of Untreated BTP

Even when persistent pain is well controlled, BTP can be devastating for you and your family or caregivers. Untreated BTP can:

- make you feel depressed and irritable.
- cause you to avoid people, even those you love.
- make you isolate yourself and prevent you from enjoying normal activities.
- keep you from getting a good night's sleep.
- interfere with doing a good job at work or even keeping it full.
- make it more difficult to relieve your persistent pain.
- increase the number of visits to your doctors and to emergency departments.

### Resources

for more information on pain management and BTP:

American Chronic Pain Association  
[www.theacpa.org](http://www.theacpa.org)

American Pain Society  
[www.am painsoc.org](http://www.am painsoc.org)

American Cancer Society  
[www.cancer.org](http://www.cancer.org)

Get a link  
[www.oncolink.com](http://www.oncolink.com)

### Managing Breakthrough Pain (BTP)

What you need to know about breakthrough pain.



Do you feel that your daily pain is fairly well controlled, yet still find there are times when you experience a sudden flare or pain that "breaks through"? These flares are called breakthrough pain or BTP.

### What Is BTP?

Breakthrough pain is an intense increase in pain that occurs suddenly even when pain-control medication is being used. Breakthrough pain can happen spontaneously or in relation to a specific activity. BTP can start and become severe in as little as three to five minutes and last an average of 30 to 60 minutes. Most people with BTP report that it happens three to four times a day. Sixty-four percent of people treated for chronic pain associated with cancer and seventy-four percent of people treated for other chronic pain conditions will experience BTP.



American Chronic Pain Association

P.O. Box 850, Rocklin, CA 95677  
(916) 632-0922; 800-533-3231

Web site: [www.theacpa.org](http://www.theacpa.org)

E-mail: [ACPA@pacbell.net](mailto:ACPA@pacbell.net)

Fax: (916) 632-3298

© 2004 American Chronic Pain Association. All rights reserved.  
Reproduction of this document is permitted by permission of the American Chronic Pain Association.



American Chronic Pain Association



### Understanding Your BTP

The first step to coping with your BTP is to learn more about it. Keep a log of when it occurs and what you were doing at the time. Note your stress levels and state of mind, as well.

Review your log every week to see if there are certain times of the day that BTP occurs. You also may find that certain activities or feelings trigger your BTP. When you know more about what causes BTP, you can begin to manage it better.

Always take your log with you to your doctor appointments. Talk to your health care provider about your concerns and fears. When your health care provider understands your issues, he or she can work with you to find a solution.

### Getting the Right Treatment for Your Pain

The key to relieving breakthrough pain is to learn why these flares of pain are different from the pain you feel all day. You can:

- Talk with your doctor about medication that might help during BTP episodes. When using pain medications for BTP, the goal should always be improved pain relief and improved function.
- Ask your doctor about biofeedback training or stress management classes. Reducing stress may help you reduce the number of episodes of BTP you have or lessen their severity.
- Understand what activities create BTP. Then you can pace your activities so you don't push yourself beyond your limits.
- Learn about the pain management skills offered through the ACPA.
- Involve your family in your treatment plans.

*For many people with moderate-to-severe chronic pain who also experience BTP, a logical treatment plan includes a pain medicine that can be taken in regular doses around the clock to treat persistent pain, plus a short-acting medicine to take when you need to relieve the rapid onset of a BTP flare. In addition, you can get involved in managing your pain by practicing relaxation techniques, setting more realistic goals, pacing yourself, and asking for help when you need it to avoid triggering flares.*

*Managing BTP is a team effort. Work with your health care team to find the overall treatment plan that's right for you.*

### Prepare for Your Next Doctor Visit

- Note if there is a regular time during the day when your persistent pain medicine doesn't last until you can take another dose.
- List methods of pain relief you have tried, such as exercise, meditation, stress reduction, etc., and how they work.
- List all the pain medicine you are taking, including over the counter medications, vitamins, and herbal supplements. Include a list of all medications or other treatments you take for any other reason, as well.
- List treatable conditions (persistent cough, nausea, constipation, etc.) that you feel might be a source of BTP.
- Take your pain log with you.
- Talk about activities your pain interferes with or the impact it has on your life.

*Like all pain, BTP can be managed but you must play an active role in the recovery process by becoming part of the treatment team. Working with your health care team, taking part in a peer support group, and learning as much as you can will help you make the transition from patient to person.*

